

The Entropy of Square-Free Words

MICHAEL BAAKE¹, VEIT ELSE^{1,2} AND UWE GRIMM^{3,†}

¹ Institut für Theoretische Physik, Universität Tübingen,
Auf der Morgenstelle 14, 72076 Tübingen, Germany

² Laboratory of Atomic and Solid State Physics,
Cornell University, Ithaca, NY 14853-2501, USA

³ Instituut voor Theoretische Fysica, Universiteit van Amsterdam,
Valckenierstraat 65, 1018 XE Amsterdam, The Netherlands

Abstract

Finite alphabets of at least three letters permit the construction of square-free words of infinite length. We show that the entropy density is strictly positive and derive reasonable lower and upper bounds. Finally, we present an approximate formula which is asymptotically exact with rapid convergence in the number of letters.

Résumé

Il est possible de construire des mots de longueur infinie sans carré sur un alphabet ayant au moins trois lettres. Nous démontrons que l'entropie du langage des mots sans carré sur un tel alphabet est strictement positive et l'encadrons par des bornes inférieure et supérieure raisonnables. Enfin, nous donnons pour l'entropie une expression approchée qui est asymptotiquement correcte et converge rapidement lorsque le nombre de lettres de l'alphabet tend vers l'infini.

Key Words: Combinatorics on words, Entropy, Substitution rules, Random tilings

1 Introduction

As is well-known, deterministic rules (like substitution rules etc.) can only result in tilings or discrete structures with vanishing *entropy density* [1] – the famous Penrose tiling is an example of this phenomenon. This tiling and many other ones appear in the description of so-called *quasicrystals* [2] for very good reason: Locally finite discrete structures (such as tilings with only finitely many prototiles) provide useful cell models for the description of the ordered state, not only crystalline but also quasi-crystalline [3]. In the latter case, one is

[†] Address after April 1996: Institut für Physik, Technische Universität, D-09130 Chemnitz, Germany

particularly interested in models with *finite* (i.e., non-vanishing) entropy density, as this is one possible mechanism to explain non-periodic order through entropic stabilization. In order to combine long-range order with a decent amount of randomness, so-called *random tiling models* have been studied in quite some detail [4], and there exists a reasonable qualitative understanding.

In view of these remarks it is clear that exactly solvable random tiling models are of interest. The standard case in one dimension is trivial as it is essentially equivalent to a Bernoulli scheme (cf. Ref. [5]). This can hardly be called ordered in any sense. In two dimensions, we know of only a few solved cases (such as the Fisher-Kasteleyn domino model (see app. E in Ref. [6]), the hexagonal random tiling model [4] or the square-triangle random tiling model [7]), and many attempts to improve this situation have failed so far, and the situation in higher dimensions is even worse. So the question arises whether one can find other examples in 1D that are more restrictive than the Bernoulli scheme but still provide reasonable toy models of (partially) ordered states. Here one can scan the vast number of examples of automata and other sequences [8], but hardly any of them yield an interesting model with positive entropy density.

One interesting class, however, is provided by infinite words in a finite alphabet that avoid the repetition of certain patterns. The simplest such case is the ensemble of *square-free words*, first studied by Thue [9, 10] and later on reinvestigated many times, see e.g. Refs. [11]–[22]. In particular, the combinatorial problems in the treatment of these systems are very interesting. It turned out in a series of independent articles [17, 18, 19] that the entropy density of square-free words in three letters is *positive*. It was conjectured that the existing upper bounds were much closer to the actual value of the entropy density than the lower bounds, and that it is close to 0.3. We will see later that the numerical value is in fact about 0.263719.

In this article, we will summarize some of the properties of the ensemble of square-free words in an alphabet $\mathcal{A}$ with finitely many letters, x say. (For a general background, we refer to Ref. [14], although we shall use slightly different notation here as a compromise between mathematical and physical literature.) We will concentrate on the case $x = 3$ for a while before we treat the general case. We present various rigorous results but also include new numerical calculations which finally guide us to the conjecture of an asymptotically correct formula for the entropy density of square-free words in x letters which turns out to be amazingly accurate already for small x .

2 Basic setup and inequalities

Let $A = \{a_1, a_2, \dots, a_x\}$ be a finite alphabet with x different letters (so $x \in \mathbb{N}$ is an integer). Then, by $\mathcal{A} = A^{\mathbb{N}_0}$, we denote the set of all words of finite length in the letters of A . This is a monoid with concatenation of words as operation and the empty word as neutral element [14]. Later on, we will restrict ourselves to subsets of $\mathcal{A}$ which are more interesting. If we

write $\ell(w)$ for the length of a word (thus $\ell(w) \in \mathbb{N}_0$ for all $w \in \mathcal{A}$), we can define (finite) subsets of $\mathcal{A}$ by

$$\mathcal{A}_n := \{w \in \mathcal{A} \mid \ell(w) = n\}. \quad (2.1)$$

Here, $\mathcal{A}_0$ consists only of the empty word and

$$\mathcal{A} = \mathcal{A}_\infty = \bigcup_{n=0}^{\infty} \mathcal{A}_n \quad (2.2)$$

which is sometimes also called the dictionary of the trivial language (we have put no rules yet, so the above language simply consists of all finite words in the alphabet A). We will generalize this in a moment, but consider only situations where the number of words of length n behaves in such a way that its logarithm divided by n defines a sequence that converges.

This is obviously so in the above case, where we have

$$|\mathcal{A}_n| = x^n. \quad (2.3)$$

Now, we can define the entropy $s = s(x)$ (it is actually an entropy density) through

$$s(x) = \lim_{n \rightarrow \infty} \frac{\log(|\mathcal{A}_n|)}{n}. \quad (2.4)$$

In our present case, we have of course

$$s(x) = \log(x) \quad (2.5)$$

which is a measure for the growth rate of $|\mathcal{A}_n|$ in n : $|\mathcal{A}_n| = \exp(n \cdot s(x))$.

Now, let us introduce the concept of repeat-free or square-free words. A word w is called *square-free* if neither w nor any substring of it is a square, otherwise w is said to contain a square. So, in an alphabet with two letters, $A = \{a, b\}$ say, a , b , ab , ba , aba and bab are square-free while all other words in these two letters are not. Consequently, as there are only finitely many square-free words in two letters, the corresponding entropy is zero. To make this more precise let us define

$$\mathcal{A}_n = \mathcal{A}_n^+ \cup \mathcal{A}_n^- \quad (2.6)$$

where $\mathcal{A}_n^-$ ($\mathcal{A}_n^+$) denotes the subset of all square-free words (square-containing words) of length n , and the right hand side of (2.6) is clearly the union of two disjoint sets. Consequently,

$$|\mathcal{A}_n^+| + |\mathcal{A}_n^-| = |\mathcal{A}_n| = x^n \quad (2.7)$$

and we introduce the abbreviation

$$\omega_n^\pm := |\mathcal{A}_n^\pm| \quad (2.8)$$

for convenience. There is always an implicit dependence on x , the (finite) number of letters of the alphabet A , but we will often suppress it when it is not needed.

One can now derive several properties of these numbers. It is clear from Eq. (2.6) that we have

$$\omega_n^+(x) + \omega_n^-(x) = x^n \quad (2.9)$$

and also the initial conditions

$$\omega_0^-(x) = 1, \omega_1^-(x) = x, \omega_0^+(x) = \omega_1^+(x) = 0. \quad (2.10)$$

Observe also that $\omega_2^+(x) = x$, as there are precisely x possibilities for words of type aa etc. If now, for x fixed, $\omega_n^-(x) = 0$ for some n , we must have $\omega_{n+m}^+(x) = x^{n+m}$ for all $m \geq 0$. This is so because no continuation of a square-containing word can become square-free and we can then use Eq. (2.9). This situation occurs with $x = 2$, but not with any larger x . We can strengthen this type of argument to obtain, for $n > 0$,

$$\omega_{n+1}^+(x) \geq x \cdot \omega_n^+(x) + \omega_n^-(x). \quad (2.11)$$

This is so because every word of $\mathcal{A}_n^+$ becomes, by adding one arbitrary letter of the alphabet (x possibilities), a word of $\mathcal{A}_{n+1}^+$, while every word of $\mathcal{A}_n^-$ can be made to one of $\mathcal{A}_{n+1}^+$ at least by repeating the last letter of it.

Now, with the initial conditions (2.10), repeated application of inequality (2.11) shows ($n > 1$)

$$\omega_n^+(x) \geq x^{n-1}. \quad (2.12)$$

It is possible to define the entropy s^+ of the dictionary of square-containing words, $\mathcal{A}_\infty^+$, which exists as an ordinary limit. We obtain

Proposition 1 *The entropy of $\mathcal{A}_\infty^+$ equals that of $\mathcal{A}_\infty$: $s^+(x) = s(x) = \log(x)$.*

The proof is a direct application of inequality (2.12):

$$s^+(x) = \lim_{n \rightarrow \infty} \frac{\log(\omega_n^+(x))}{n} \geq \log(x) \cdot \lim_{n \rightarrow \infty} \frac{n-1}{n} = \log(x).$$

On the other hand, we have $s^+(x) \leq s(x) = \log(x)$, from which the statement follows. $\square$

For the other entropy, $s^-(x)$, we have to prove existence as an ordinary limit first. We do that in a slightly more general form, following an argument given by Pleasants [23].

Lemma 1 *Let $a(n)$ be a sequence of positive integers with $a(m+n) \leq a(m) \cdot a(n)$. Then the sequence defined through $h(n) := \log(a(n))/n$ is convergent.*

PROOF: Let us take two integers $N \geq n$ related by $N = qn + r$ with $0 \leq r < n$. We then have, with $x := a(1)$,

$$a(N) \leq (a(n))^q \cdot a(r) \leq (a(n))^q \cdot x^r \quad (2.13)$$

and, consequently,

$$\frac{\log(a(N))}{N} \leq \frac{q \cdot \log(a(n))}{N} + \frac{r \cdot \log(x)}{N} < \frac{\log(a(n))}{n} + \frac{\log(x)}{q}. \quad (2.14)$$

This means $h(N) < h(n) + \frac{1}{q} \log(x)$ for any $N = qn + r$ with $0 \leq r < n$. But then, we can fix n and take the limit $N \rightarrow \infty$ which also implies $q \rightarrow \infty$. This gives $\limsup_{N \rightarrow \infty} h(N) \leq h(n)$. The last equation is valid for all $n \in \mathbb{N}$, so we also have

$$\limsup_{N \rightarrow \infty} h(N) \leq \liminf_{n \rightarrow \infty} h(n) \quad (2.15)$$

which means that both must be equal and the limit exists. $\square$

If $a(n)$ is only a sequence of non-negative integers, but still with $a(m+n) \leq a(m)a(n)$, one either has $a(n) > 0$ for all n or, if $a(n_0) = 0$ for some n_0 , one has $a(n) = 0$ also for all $n > n_0$ due to submultiplicativity. In the latter case, we follow the usual convention and define $h(n) = 0$ which results in $\lim_{n \rightarrow \infty} h(n) = 0$, i.e. vanishing entropy.

It is clear that the above Lemma is not the most general formulation of the statement, but it is sufficient for our needs. The language of square-free words is *subword-closed*, i.e., no new substring of length n can occur in any word of length $> n$ [23]. Consequently, our numbers $\omega_n^-(x)$ are such that the lemma applies. We have thus shown

Proposition 2 *The entropy $s^-(x)$ of $\mathcal{A}_\infty^-(x)$ exists as a limit.*

We so far only know the trivial inequality

$$0 \leq s^-(x) \leq \log(x) \quad (2.16)$$

with $s^-(x) = 0$ for $x = 1$ and $x = 2$. Since we have at most $x - 1$ possibilities to make a square-free word of length n into one of length $n + 1$ by adding a letter, we also have ($n \geq 1$)

$$\omega_{n+1}^-(x) \leq (x - 1) \cdot \omega_n^-(x) \quad (2.17)$$

from which we can improve the upper bound of (2.16) to

$$s^-(x) \leq \log(x - 1) . \quad (2.18)$$

Of course, we can further improve the upper bound by considering the possibilities of appending (square-free) words which consist of more than one letter. In practice, this amounts to actually counting the number of square-free words of a certain length. One obtains

$$\omega_{n+k}^-(x) \leq \frac{\omega_{k+j}^-(x)}{\omega_j^-(x)} \omega_n^-(x) \quad (2.19)$$

where $j \in \{0, 1, 2\}$ and $n \geq j$ and $k \geq 0$ are arbitrary. This expression follows by extending square-free words of length n with an overlapping square-free word of length $k + j$. By restricting the length of the overlap to $j \leq 2$ all possibilities appear with the same frequency (by symmetry). For the entropy, this yields upper bounds

$$s^-(x) \leq \frac{1}{k} \left(\log(\omega_{k+j}^-(x)) - \log(\omega_j^-(x)) \right) \quad (2.20)$$

which clearly gives the strongest bound for $j = 2$, with $\omega_2^-(x) = x(x - 1)$.

Now, a more interesting point is the question for a lower bound of the entropy $s^-(x)$. We know it is zero for $x = 1$ and $x = 2$. As we will show, it is strictly positive for other x , i.e. for $x > 2$. The case $x = 3$ will play a special role, but let us first give some simple results. Here, we rely on the well-known fact that there exists at least one square-free word of infinite length in three letters, compare Ref. [14] and references therein. But then, we can directly show

Theorem 1 For $x \geq 3$, the entropy of $\mathcal{A}_\infty^-$ has a lower bound: $s^-(x) \geq \frac{1}{3} \log(x-2)$. In particular, the entropy is strictly positive for $x > 3$.

PROOF: Let w be a square-free word of infinite length in 3 letters, $\{a, b, c\}$ say, which we know to exist from Refs. [9, 14]. Let w_n be the subword made from the first n letters of w which is also square-free and contains m_a, m_b, m_c letters of type a, b, c , respectively, where $m_a + m_b + m_c = n$. Now, let $\{d_1, d_2, \dots, d_p\}$ be p new letters, $p > 0$. If we fix all b 's and c 's in w_n , we have $(p+1)^{m_a}$ possibilities to make square-free words of length n in four letters by successive replacement of any a in w by an element of $\{a, d_1, \dots, d_p\}$. We can analogously proceed for the other two letters, b and c , through fixing a, c and a, b , respectively. In this setup, we have $x = p + 3$, and we can conclude (for $x \geq 3$)

$$\omega_n^-(x) \geq (x-2)^{m_a} + (x-2)^{m_b} + (x-2)^{m_c} \geq 3 \cdot (x-2)^{n/3} \quad (2.21)$$

where the second inequality is a standard result from calculus. But from this, we immediately get the inequality

$$s^-(x) \geq \frac{1}{3} \log(x-2) + \lim_{n \rightarrow \infty} \frac{3}{n} = \frac{1}{3} \log(x-2) \quad (2.22)$$

from which the statement follows. $\square$

We cannot gain anything about $\varepsilon := s^-(3)$ this way, although, as we will see, precisely this ε is important. Nevertheless, we can do better than (2.22). In the above argument, we started from an infinite word w in three letters. Instead, we can also apply the same type of argument for the step from x to $x+1$ letters: fixing $x-1$ letters, we stay with two possibilities for the replacement of every occurrence of the remaining letter, and this can be done in x different ways. As it applies essentially to every word separately, the number of possibilities behaves almost multiplicatively, i.e. it grows like

$$\omega_n^-(x+1) \sim x \cdot 2^{n/x} \cdot \omega_n^-(x). \quad (2.23)$$

We cannot write $\geq$ instead of $\sim$ here, as one does in fact multiply count several words.

To avoid this, we have to discard all words of $\mathcal{A}_n^-(x)$ that do not contain all letters. Let us denote the number of square-free words of length n in *exactly* x letters by $\psi_n(x)$. Clearly, $\psi_n(0) = \delta_{n,0}$, $\psi_n(1) = \delta_{n,1}$ and $\psi_n(2) = 2(\delta_{n,2} + \delta_{n,3})$. Also, $\psi_n(3) = \omega_n^-(3) > 0$ for all $n > 3$. Furthermore, we have

$$\omega_n^-(x) = \sum_{k=0}^x \binom{x}{k} \psi_n(k) \quad (2.24)$$

and also, obviously, $\omega_n^-(x) \geq \psi_n(x)$ for all n, x . For $x > n$, one has $\psi_n(x) = 0$, and $\psi_x(x) = x!$, the number of permutations of x symbols. By simple counting, one finds $\psi_{x+1}(x) = x!x(x-1)/2$. A little less obvious is the inversion of Eq. (2.24) in the form

$$\psi_n(x) = \sum_{k=0}^x (-1)^{x-k} \binom{x}{k} \omega_n^-(k). \quad (2.25)$$

The number $\psi_n(x)$ is a multiple of $x!$ since any permutation of the x letters transforms a square-free word in exactly x letters into another one of this type. The advantage of the new numbers is that we can go from $\psi_n(x)$ to $\psi_n(x+1)$ without any double counting, i.e., two different square-free words of length n with all x letters in them give automatically two disjoint sets of square-free words of length n in $x+1$ letters by the above procedure. If we observe that we must introduce at least one new letter, we see that, for $x \geq 1$ and $n \geq x+1$, we get the inequality

$$\psi_n(x+1) \geq \frac{x}{2} \cdot 2^{n/x} \cdot \psi_n(x). \quad (2.26)$$

This is helpful because we have:

Lemma 2 *Let $x \geq 3$ be fixed. Then, the sequences $\psi_n(x)$ and $\omega_n^-(x)$ have the same exponential growth in n :*

$$\lim_{n \rightarrow \infty} \frac{\log(\psi_n(x))}{n} = \lim_{n \rightarrow \infty} \frac{\log(\omega_n^-(x))}{n}.$$

PROOF: Iterating Eq. (2.26) one obtains (for $0 < k < x$ and $n \geq x$) the inequality

$$\psi_n(k) \leq \psi_n(x) \cdot \left(\frac{2^{x-k}}{2^{n(\frac{1}{k} + \dots + \frac{1}{x-1})}} \cdot \frac{(k-1)!}{(x-1)!} \right). \quad (2.27)$$

Consequently, we obtain, from Eq. (2.24), the inequality

$$\omega_n^-(x) \leq \psi_n(x) \left[1 + \sum_{k=1}^{x-1} \frac{x}{k} \cdot \frac{2^{x-k}}{(x-k)!} \cdot \frac{1}{2^{n(\frac{1}{k} + \dots + \frac{1}{x-1})}} \right]. \quad (2.28)$$

Since every single term under the sum is certainly not bigger than $2x$, we finally get

$$\psi_n(x) \leq \omega_n^-(x) \leq (1 + 2x(x-1)) \cdot \psi_n(x) \leq (1 + 2x(x-1)) \cdot \omega_n^-(x) \quad (2.29)$$

from which the statement easily follows. $\square$

From this Lemma and from Eq. (2.26) we see that the entropic contributions are additive:

$$s^-(x+1) \geq s^-(x) + \frac{\log(2)}{x} \quad (2.30)$$

which is valid for $x \geq 3$. In fact, repeating the argument of (2.30), we immediately arrive at

Theorem 2 *Let $x > 3$. The entropy of the dictionary $\mathcal{A}_\infty^-(x)$ fulfills:*

$$\varepsilon + \log(2) \cdot \left(\frac{1}{3} + \frac{1}{4} + \dots + \frac{1}{x-1} \right) \leq s^-(x) \leq \log(x-1).$$

Here, and in what follows, we will always use ε for $s^-(3)$. Of course, one can now use the formula

$$\sigma_m := 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{m} - \log(m) \xrightarrow{m \rightarrow \infty} \gamma \simeq 0.5771 \dots \quad (2.31)$$

for Euler's constant and the fact that the sequence $(\sigma_m)_{m \in \mathbf{N}}$ is strictly decreasing to simplify the inequality of Thm. (2) for larger values of x , while for small values it is better to stick to the finite sum.

Again, we have seen that ε plays a special role. In fact, although this does not follow from any simple inequality of the above type, ε is strictly positive [17, 18, 19]. Brandenburg [18] shows that

$$\omega_{22n}^-(3) \geq 2^n \omega_n^-(3) \quad (2.32)$$

and concludes from this that

$$\omega_n^-(3) \geq 6 \cdot 2^{n/22}. \quad (2.33)$$

In fact, using the *existence* of the limit (which we have shown above), we can improve the argument slightly (cf. Ref. [17, 19]) to obtain a strict lower bound for the entropy of square-free words on three letters:

Theorem 3 *The entropy $\varepsilon = s^-(3)$ is strictly positive:*

$$\varepsilon \geq \frac{1}{21} \log(2) \simeq 0.033007 \dots \quad (2.34)$$

Since this kind of result has been described several times already [17, 18, 19], we shall not repeat the proof. The idea behind it is the following: one tries to find a set of substitution rules which map square-free words into square-free words of increased length and simultaneously allow some free choice to do so in each step – which accounts for the exponential growth. It is not clear whether one can significantly improve the lower bound in this way so as to approach our numerical estimate of

$$\varepsilon \simeq 0.263719(1). \quad (2.35)$$

3 Some results for three letters

Let us describe the case $x = 3$ in more detail. We can write the set of square-free words in three letters as a disjoint union of three subsets

$$\mathcal{A}_n^- = \mathcal{A}_n^{(0)} \cup \mathcal{A}_n^{(1)} \cup \mathcal{A}_n^{(2)}. \quad (3.1)$$

Here, $\mathcal{A}_n^{(0)}$ denotes the set of what we call *stop-words*. These are characterized by the property that by appending *any* letter of the alphabet A one obtains a square-containing word. In the same spirit, $\mathcal{A}_n^{(1)}$ and $\mathcal{A}_n^{(2)}$ are defined as the sets of square-free words which allow, respectively, one and two extensions to a square-free word of length $n+1$. Introducing the notation

$$\omega_n^{(k)} := |\mathcal{A}_n^{(k)}|, \quad k \in \{0, 1, 2\}, \quad (3.2)$$

this implies the relation

$$\omega_{n+1}^- = \sum_{k=0}^2 \omega_{n+1}^{(k)} = \sum_{k=0}^2 k \cdot \omega_n^{(k)} = \omega_n^{(1)} + 2 \cdot \omega_n^{(2)}. \quad (3.3)$$

Hence the growth rate is given by

$$\frac{\omega_{n+1}^-}{\omega_n^-} = 1 + \frac{\omega_n^{(2)} - \omega_n^{(0)}}{\omega_n^-}. \quad (3.4)$$

Since the left-hand side converges to a finite value in the limit $n \rightarrow \infty$, so does the right-hand side, and by the positivity of the entropy we see that convergence of the sequence $\omega_n^{(0)}/\omega_n^-$ would imply the convergence of the ratio $\omega_n^{(2)}/\omega_n^-$ to a finite non-zero limit.

In order to gain some insight into the actual behaviour of these quantities we used the computer to investigate square-free words in three letters for lengths up to 90. The results are presented in Tables 1 and 2.

In Table 1, we list the number of square-free words ω_n^- , approximants to the entropy obtained from the ratio of successive values, and upper limits to the entropy using Eq. (2.20). From these, we extract

$$\varepsilon \simeq 0.263719(1) \quad (3.5)$$

as the approximate value of the entropy of square-free words on three letters, where the figure in parentheses denotes the estimated uncertainty in the last digit.

It is striking that the logarithm of the ratio obviously approaches the limit value much faster than the value in the last column. This in fact suggests that the asymptotic behaviour looks as follows

$$\log(\omega_n^-) \sim \varepsilon \cdot n + a + o(n^{-1}) \quad (n \rightarrow \infty) \quad (3.6)$$

where the constant term can be estimated as $a \simeq 2.5438965$. It is interesting that the next order seems to fall so quickly, which also indicates that no logarithmic corrections are present.

Table 2 contains the values of $\omega_n^{(k)}$ ($k = 0, 1, 2$) and their ratios with ω_n^- . Apparently, the ratios converge as $n \rightarrow \infty$, which means that all three subsets have in fact the same entropy as the set of square-free words itself. For the ratios, we estimate

$$\frac{\omega_n^{(0)}}{\omega_n^-} \rightsquigarrow 0.036837, \quad \frac{\omega_n^{(1)}}{\omega_n^-} \rightsquigarrow 0.624564, \quad \frac{\omega_n^{(2)}}{\omega_n^-} \rightsquigarrow 0.338599, \quad (3.7)$$

in the limit $n \rightarrow \infty$, where the uncertainty is about one figure in the last digit.

We note that the stop-words still show an interesting structure if one looks at the three lengths of squares that one obtains on appending the three different letters. Apparently, stop-words with certain fixed sets of periods still occur with the same entropy density, whereas other periods are limited to “symmetric” stop-words that cannot be extended in any direction and therefore show up in finitely many stop-words only. As an example for this, we mention the shortest stop-words (*abacaba* and those obtained from permutation of letters, where a , b and c denote the three letters) which result in squares of lengths 1, 2 and 4. Here, it is easy to see that these are the only words with this property.

Table 1: Number of square-free words in three letters; estimates and upper limits for their entropy.

n	ω_n^-	$\log(\omega_n^-/\omega_{n-1}^-)$	$\frac{\log(\omega_n^-/6)}{n-2}$
1	3		
2	6	0.69314718	
3	12	0.69314718	0.69314718
4	18	0.40546511	0.54930615
5	30	0.51082562	0.53647929
6	42	0.33647224	0.48647752
7	60	0.35667494	0.46051702
8	78	0.26236426	0.42749155
9	108	0.32542240	0.41291025
10	144	0.28768207	0.39725673
11	204	0.34830669	0.39181784
12	264	0.25782911	0.37841898
13	342	0.25886163	0.36755010
14	456	0.28768207	0.36089444
15	618	0.30399565	0.35651761
16	798	0.25562014	0.34931064
17	1044	0.26870617	0.34393701
18	1392	0.28768207	0.34042108
19	1830	0.27357440	0.33648893
20	2388	0.26614023	0.33258069
21	3180	0.28642500	0.33015144
22	4146	0.26526282	0.32690698
23	5418	0.26758273	0.32408205
24	7032	0.26074442	0.32120302
25	9198	0.26851491	0.31891227
26	11892	0.25687984	0.31632757
27	15486	0.26407048	0.31423730
28	20220	0.26673582	0.31241032
29	26424	0.26760047	0.31075069
30	34422	0.26442321	0.30909613
31	44862	0.26489522	0.30757198
32	58446	0.26451214	0.30613664
33	76122	0.26423407	0.30478492
34	99276	0.26556653	0.30355936
35	129516	0.26590058	0.30241820
36	168546	0.26340428	0.30127072
37	219516	0.26421641	0.30021203
38	285750	0.26369218	0.29919758
39	372204	0.26432479	0.29825506
40	484446	0.26356388	0.29734215
41	630666	0.26377043	0.29648134
42	821154	0.26393426	0.29566768
43	1069512	0.26424708	0.29490131
44	1392270	0.26373304	0.29415920
45	1812876	0.26397903	0.29345734

Table 1: (continued)

n	ω_n^-	$\log(\omega_n^-/\omega_{n-1}^-)$	$\frac{\log(\omega_n^-/6)}{n-2}$
46	2359710	0.26362420	0.29277931
47	3072486	0.26394828	0.29213862
48	4000002	0.26380786	0.29152274
49	5207706	0.26384459	0.29093384
50	6778926	0.26367923	0.29036604
51	8824956	0.26376493	0.28982316
52	11488392	0.26375352	0.28930176
53	14956584	0.26381447	0.28880201
54	19470384	0.26374294	0.28832011
55	25346550	0.26374808	0.28785649
56	32996442	0.26375711	0.28741020
57	42957300	0.26380686	0.28698105
58	55921896	0.26374940	0.28656620
59	72798942	0.26374542	0.28616583
60	94766136	0.26371071	0.28577868
61	123368406	0.26376292	0.28540553
62	160596120	0.26371759	0.28504406
63	209059806	0.26372772	0.28469461
64	272143380	0.26370870	0.28435613
65	354271314	0.26373398	0.28402880
66	461181036	0.26372763	0.28371159
67	600356406	0.26373282	0.28340422
68	781520994	0.26371852	0.28310596
69	1017362166	0.26372643	0.28281671
70	1324371090	0.26372453	0.28253594
71	1724034504	0.26372949	0.28226338
72	2244278358	0.26371684	0.28199843
73	2921521164	0.26372040	0.28174100
74	3803130042	0.26372000	0.28149071
75	4950798954	0.26372455	0.28124733
76	6444761514	0.26371866	0.28101046
77	8389549680	0.26371921	0.28077991
78	10921197582	0.26371879	0.28055542
79	14216853012	0.26372246	0.28033681
80	18506985300	0.26372015	0.28012378
81	24091726728	0.26372025	0.27991614
82	31361678988	0.26371824	0.27971366
83	40825520274	0.26372065	0.27951622
84	53145145482	0.26371938	0.27932357
85	69182396616	0.26371968	0.27913558
86	90058945560	0.26371796	0.27895203
87	117235364616	0.26371917	0.27877282
88	152612592438	0.26371906	0.27859778
89	198665414208	0.26371944	0.27842676
90	258615015792	0.26371846	0.27825962

Table 2: Number and relative frequency of square-free words in three letters that allow 0, 1, and 2 extensions, respectively.

n	$\omega_n^{(0)}$	$\omega_n^{(1)}$	$\omega_n^{(2)}$	$\omega_n^{(0)}/\omega_n^-$	$\omega_n^{(1)}/\omega_n^-$	$\omega_n^{(2)}/\omega_n^-$
1	0	0	3	0.00000000	0.00000000	1.00000000
2	0	0	6	0.00000000	0.00000000	1.00000000
3	0	6	6	0.00000000	0.50000000	0.50000000
4	0	6	12	0.00000000	0.33333333	0.66666667
5	0	18	12	0.00000000	0.60000000	0.40000000
6	0	24	18	0.00000000	0.57142857	0.42857143
7	6	30	24	0.10000000	0.50000000	0.40000000
8	0	48	30	0.00000000	0.61538462	0.38461538
9	0	72	36	0.00000000	0.66666667	0.33333333
10	0	84	60	0.00000000	0.58333333	0.41666667
11	6	132	66	0.02941176	0.64705882	0.32352941
12	6	174	84	0.02272727	0.65909091	0.31818182
13	6	216	120	0.01754386	0.63157895	0.35087719
14	6	282	168	0.01315789	0.61842105	0.36842105
15	24	390	204	0.03883495	0.63106796	0.33009709
16	24	504	270	0.03007519	0.63157895	0.33834586
17	24	648	372	0.02298851	0.62068966	0.35632184
18	36	882	474	0.02586207	0.63362069	0.34051724
19	54	1164	612	0.02950820	0.63606557	0.33442623
20	54	1488	846	0.02261307	0.62311558	0.35427136
21	120	1974	1086	0.03773585	0.62075472	0.34150943
22	138	2598	1410	0.03328509	0.62662808	0.34008683
23	216	3372	1830	0.03986711	0.62236988	0.33776301
24	240	4386	2406	0.03412969	0.62372014	0.34215017
25	384	5736	3078	0.04174821	0.62361383	0.33463796
26	444	7410	4038	0.03733602	0.62310797	0.33955600
27	528	9696	5262	0.03409531	0.62611391	0.33979078
28	690	12636	6894	0.03412463	0.62492582	0.34094955
29	966	16494	8964	0.03655767	0.62420527	0.33923706
30	1236	21510	11676	0.03590727	0.62489106	0.33920167
31	1602	28074	15186	0.03570951	0.62578574	0.33850475
32	2112	36546	19788	0.03613592	0.62529514	0.33856894
33	2712	47544	25866	0.03562702	0.62457634	0.33979664
34	3522	61992	33762	0.03547685	0.62444095	0.34008220
35	4818	80850	43848	0.03720004	0.62424720	0.33855277
36	6150	105276	57120	0.03648856	0.62461287	0.33889858
37	8094	137094	74328	0.03687203	0.62452851	0.33859946
38	10452	178392	96906	0.03657743	0.62429396	0.33912861
39	13854	232254	126096	0.03722152	0.62399652	0.33878196
40	17784	302658	164004	0.03670997	0.62475075	0.33853928
41	23082	394014	213570	0.03659940	0.62475859	0.33864201
42	29970	512856	278328	0.03649742	0.62455520	0.33894738
43	39438	667878	362196	0.03687476	0.62446985	0.33865539
44	51030	869604	471636	0.03665237	0.62459437	0.33875326
45	66792	1132458	613626	0.03684312	0.62467483	0.33848206

Table 2: (continued)

n	$\omega_n^{(0)}$	$\omega_n^{(1)}$	$\omega_n^{(2)}$	$\omega_n^{(0)}/\omega_n^-$	$\omega_n^{(1)}/\omega_n^-$	$\omega_n^{(2)}/\omega_n^-$
46	86502	1473930	799278	0.03665789	0.62462336	0.33871874
47	113064	1918842	1040580	0.03679887	0.62452425	0.33867689
48	147036	2498226	1354740	0.03675898	0.62455619	0.33868483
49	191952	3252582	1763172	0.03685922	0.62457097	0.33856980
50	249390	4234116	2295420	0.03678901	0.62459983	0.33861116
51	324852	5511816	2988288	0.03681061	0.62457150	0.33861789
52	422712	7174776	3890904	0.03679471	0.62452395	0.33868134
53	550758	9341268	5064558	0.03682378	0.62455892	0.33861729
54	716454	12161310	6592620	0.03679712	0.62460555	0.33859733
55	932592	15831474	8582484	0.03679365	0.62460074	0.33860561
56	1213602	20608380	11174460	0.03677978	0.62456370	0.33865651
57	1582026	26828652	14546622	0.03682787	0.62454232	0.33862980
58	2058528	34927794	18935574	0.03681077	0.62458172	0.33860751
59	2681796	45468156	24648990	0.03683839	0.62457166	0.33858995
60	3488478	59186910	32090748	0.03681144	0.62455760	0.33863096
61	4545588	77049516	41773302	0.03684564	0.62454820	0.33860616
62	5914926	100302582	54378612	0.03683106	0.62456417	0.33860477
63	7701792	130572648	70785366	0.03684014	0.62457079	0.33858907
64	10021482	169972482	92149416	0.03682427	0.62456960	0.33860613
65	13049082	221263428	119958804	0.03683358	0.62455926	0.33860716
66	16985274	288035118	156160644	0.03682995	0.62455976	0.33861029
67	22114344	374963130	203278932	0.03683536	0.62456755	0.33859709
68	28782414	488114994	264623586	0.03682872	0.62457055	0.33860074
69	37472418	635408406	344481342	0.03683292	0.62456461	0.33860247
70	48778746	827150184	448442160	0.03683163	0.62456074	0.33860763
71	63510756	1076769138	583754610	0.03683845	0.62456357	0.33859799
72	82666266	1401703020	759909072	0.03683423	0.62456736	0.33859840
73	107616300	1824679686	989225178	0.03683571	0.62456494	0.33859935
74	140084994	2375291142	1287753906	0.03683413	0.62456217	0.33860370
75	182377848	3092080698	1676340408	0.03683806	0.62456196	0.33859998
76	237398214	4025176920	2182186380	0.03683584	0.62456569	0.33859847
77	309038124	5239825530	2840686026	0.03683608	0.62456577	0.33859815
78	402276216	6820989720	3697931646	0.03683444	0.62456426	0.33860130
79	523700664	8879319396	4813832952	0.03683661	0.62456293	0.33860046
80	681718896	11558806080	6266460324	0.03683576	0.62456450	0.33859974
81	887460042	15046854384	8157412302	0.03683671	0.62456521	0.33859808
82	1155219294	19587399114	10619060580	0.03683538	0.62456475	0.33859987
83	1503883698	25498127670	13823508906	0.03683685	0.62456345	0.33859970
84	1957680408	33192533532	17994931542	0.03683649	0.62456379	0.33859972
85	2548490760	43208866152	23425039704	0.03683727	0.62456446	0.33859827
86	3317442636	56247641232	30493861692	0.03683635	0.62456473	0.33859892
87	4318568760	73220999274	39695796582	0.03683674	0.62456409	0.33859917
88	5621734092	95316302484	51674555862	0.03683663	0.62456381	0.33859955
89	7318253526	124079305572	67267855110	0.03683708	0.62456420	0.33859872

4 Basics of an algebraic approach

It is now time to attack the square-free words with some algebra. To do so, we prefer to change the notation

$$P_n(x) := |\mathcal{A}_n^-(x)| = \omega_n^-(x) \quad (4.1)$$

because we will not talk about square-containing words any more. Clearly, $P_n(x)$ is always an integer, and we also know that, for $n \in \mathbb{N}$, $P_n(x) \leq x^n$, with equality only for $n = 0$ and $n = 1$. It is straight-forward to calculate the first cases explicitly

$$\begin{aligned} P_0(x) &= 1, \quad P_1(x) = x, \quad P_2(x) = x(x-1), \\ P_3(x) &= x(x-1)^2, \quad P_4(x) = x^2(x-1)(x-2). \end{aligned} \quad (4.2)$$

The $P_n(x)$ are polynomials in x , which can be shown by induction. If we go to $P_{n+1}(x)$ we can recursively build it from $(x-1)P_n(x)$ corrected by lower order terms that are sums of products of $P_k(x)$ with $k < n$ – hence we stay with a polynomial.

Let us consider these polynomials in more detail. If $x > 2$ and $n \geq 2$, we can fix the first two letters when we want to count the square-free letters of length n as ab say. For this we have $x(x-1)$ possibilities, each of which must have equally many square-free continuations. Similarly, let $x > 3$ and $n \geq 4$. The start can be of the form abc (where one has $x(x-1)(x-2)$ possibilities, each with equally many square-free continuations) or of the form $abac$ (again $x(x-1)(x-2)$ possibilities). The set of words obtained from these two classes of possibilities is disjoint, so we can conclude

Proposition 3 *Let $P_n(x)$ be the number of square-free words of length n in an alphabet of x letters. Then, for $n > 1$, $P_n(x)$ is a multiple of $x(x-1)$, while for $n > 3$, it is a multiple of $x(x-1)(x-2)$. Furthermore, $P_n(x)$ is a polynomial in x of order n , with integer coefficients and leading coefficient 1.*

There is an alternative way to see that the last statement of the lemma is correct. Clearly, the number of all words of length n in x letters is just x^n , and from this we have to subtract the number of words which contain at least one square. Necessarily, demanding that a word contains a square of a certain length (and no square of shorter length) means that one fixes a number of letters to coincide (and certain others to be different), thus all the corresponding terms are of lower order in x . Also, they all have integer coefficients since these are nothing but combinatorial factors. Furthermore, as $P_n(x) \leq x^n$, the coefficient of the leading term of $P_n(x) - x^n$ has to be negative.

So far, we did not manage to find the generating function for the polynomials $P_n(x)$ – and there are the usual indications that this might be a very difficult task: quite probably it will be one of those functions that are not analytically continuable beyond its circle of convergence. We assume this due to various unsuccessful attempts to find generating functions for the numbers $\omega_n^-(x)$ by means of standard algebraic program packages – although

it is certainly not conclusive. So, we have tried to find a reasonable approximation scheme. Here, we observe, from the explicit counting and the determination of the $P_n(x)$ up to $n = 15$, that the recurrence of the polynomials, for $n > 2$, look as follows

$$P_{n+1}(x) = (x-1) \cdot P_n(x) - P_{n-1}(x) + R_{n-2}(x) . \quad (4.3)$$

Here, the remainder $R_{n-2}(x)$ is a polynomial in x of *at most* degree $n-2$, but usually of considerably smaller degree (though we do not know how to quantify this at the moment). The first two terms on the right hand side of (4.3) are clear: one has $x-1$ possibilities to extend a square-free word of length n into one of length $n+1$ without a square of length 2 at the end. From these (if $n > 2$) essentially $P_{n-1}(x)$ words have to be subtracted because they contain a square of length 4 at the end. Further restrictions reach deeper into the word, as can be seen from the structure of the stop-words.

Consequently, at least for large x , one would expect a reasonable approximation by simply neglecting the lower order terms. So, let us consider polynomials $Q_n(x)$ defined through

$$Q_{n+1}(x) = (x-1) \cdot Q_n(x) - Q_{n-1}(x) \quad (4.4)$$

with initial conditions $Q_0(x) \equiv 1$ and $Q_{-1}(x) \equiv 0$. In this case, the generating function can easily be calculated

$$F(x, t) := \sum_{n=0}^{\infty} Q_n(x) \cdot t^n = \frac{1}{t^2 - (x-1)t + 1} \quad (4.5)$$

by applying the recurrence relation (4.4) and observing the initial conditions properly.

For a given x , the growth rate of the coefficients (which we know to converge) is given by the inverse of the radius of convergence ϱ of the generating function wherefore we obtain the simple formula

$$\tilde{s}(x) = \log \left(\frac{(x-1) + \sqrt{(x-1)^2 - 4}}{2} \right) . \quad (4.6)$$

Although the first two terms in the asymptotic expansion of (4.6) match those of the upper bound, $\log(x-1) \sim \log(x) - x^{-1}$, for finite x it gives a much better approximation to the true value of the entropy (see Table 3) except for $x = 3$, where $\tilde{s}(3) = 0$. The lower bounds are those due to Proposition 1 with $\varepsilon = \log(2)/21$ while the estimate is obtained from counting square-free words in x letters up to length $n_{\max}$ (by extrapolating the logarithm of the successive ratios). The error is roughly 1 figure in the last digit. The upper bound is again strict and was calculated from Eq. (2.20) as

$$s^-(x) \leq \frac{1}{n_{\max} - 2} \log \left(\frac{\omega_{n_{\max}}^-(x)}{x(x-1)} \right) . \quad (4.7)$$

The convergence of $\tilde{s}(x)$ is rather quick (see Table 3), and the above arguments indicate that $\tilde{s}(x)$ is asymptotically exact – a property that deserves further investigation.

Table 3: Bounds and estimates for the entropy of square-free words in x letters.

x	$n_{\max}$	lower bound	estimate	upper bound	$\log(x-1)$	$\tilde{s}(x)$
3	90	0.03300701	0.263719	0.27825962	0.69314718	0.00000000
4	26	0.26405607	0.96375	0.97319304	1.09861229	0.96242365
5	21	0.43734286	1.317089	1.32225469	1.38629436	1.31695790
6	18	0.57597230	1.56682	1.57028618	1.60943791	1.56679924
7	16	0.69149683	1.76275	1.76530829	1.79175947	1.76274717
8	16	0.79051786	1.924850	1.92663981	1.94591015	1.92484730
9	15	0.87716125	2.063438	2.06486642	2.07944154	2.06343707
10	14	0.95417761	2.184644	2.18583786	2.19722458	2.18464379
11	12	1.02349232	2.29243	2.29357100	2.30258509	2.29243167
12	12	1.08650570	2.38953	2.39045454	2.39789527	2.38952643

5 Concluding remarks

In this article, we have discussed various aspects of the ensemble of square-free words in a finite alphabet with x letters, with some emphasis on the entropy density in the thermodynamic limit. Though we could give various rigorous bounds for the entropy (which is, in particular, strictly positive for $x > 2$), we were neither able to solve the problem analytically nor able to construct an exhaustive lower bound (while this is easy for the upper bound). Nevertheless, an approximate generating function was given which results in an entropy estimate that is asymptotically exact and astonishingly accurate already for small $x > 3$.

Several questions remain open. Although standard criteria point to non-solvability of the problem (in the sense that the generating function has the circle of convergence as its analyticity domain), this needs further investigation. Here, a better understanding of the lower bound would help because it would shed more light onto possible methods to essentially exhaust the square-free words, at least w.r.t. their exponential growth. We hope to report on further findings in this direction in the near future.

Acknowledgements

It is a pleasure to thank J.-P. Allouche and J. Cassaigne for discussions and various useful comments. M. B. would like to thank B. Iochum and the CPT Luminy for hospitality and financial support during a stay in spring 1995 where part of this work was done. V. E. thanks the Alexander von Humboldt Foundation for support. U. G. gratefully acknowledges financial support of the Samenwerkingverband FOM/SMC Mathematische Fysica.

References

- [1] V. Berthé, “Entropy in deterministic and random systems”, in: F. Axel and D. Gratias (eds.), *Beyond Quasicrystals*, Springer, Berlin (1995), 441–463.
- [2] C. Janot, *Quasicrystals – a Primer*, 2nd ed., Clarendon Press, Oxford (1994).
- [3] H.-U. Nissen and C. Beeli, “Electron microscopy of quasicrystals and the validity of the tiling approach”, *Int. J. Mod. Phys.* **B7** (1993) 1387–1413.
- [4] C. L. Henley, “Random Tiling Models”, in: D. P. DiVincenzo and P. J. Steinhardt (eds.), *Quasicrystals – the State of the Art*, World Scientific, Singapore (1991), 429–524.
- [5] K. Petersen, *Ergodic Theory*, Cambridge University Press, Cambridge (1983).
- [6] C. J. Thompson, *Mathematical Statistical Mechanics*, Princeton University Press, Princeton (1979).
- [7] P. A. Kalugin, “The square-triangle random-tiling model in the thermodynamic limit”, *J. Phys.* **A27** (1994) 3599–3614.
- [8] J.-P. Allouche and M. Mendès France, “Automata and automatic sequences”, in: F. Axel and D. Gratias (eds.), *Beyond Quasicrystals*, Springer, Berlin (1995), 293–367.
- [9] A. Thue, “Über unendliche Zeichenreihen”, *Norske Vid. Selsk. Skr. I, Mat. Nat. Kl. Christiana* **7** (1906) 1-22; reprinted in: T. Nagell, A. Selberg, S. Selberg and K. Thalberg (eds.), *Selected Mathematical Papers of Axel Thue*, Universitetsforlaget, Oslo (1977), 139–158.
- [10] A. Thue, “Über die gegenseitige Lage gleicher Teile gewisser Zeichenreihen”, *Norske Vid. Selsk. Skr. I, Mat. Nat. Kl. Christiana* **1** (1912) 1-67; reprinted in: T. Nagell, A. Selberg, S. Selberg and K. Thalberg (eds.), *Selected Mathematical Papers of Axel Thue*, Universitetsforlaget, Oslo (1977), 413–477.
- [11] P. A. B. Pleasants, “Nonrepetitive sequences”, *Proc. Cambr. Philos. Soc.* **68** (1970) 267–274.
- [12] D. R. Bean, A. Ehrenfeucht and G. F. McNulty, “Avoidable patterns in strings of symbols”, *Pacific J. Math.* **85** (1979) 261–294.
- [13] M. Crochemore, “Sharp characterizations of squarefree morphisms”, *Theor. Comp. Sci.* **18** (1982) 221–226.
- [14] J. Berstel and C. Reutenauer, “Square-Free Words and Idempotent Semigroups”, in: *Combinatorics on Words*, ed. M. Lothaire, Encyclopedia of mathematics and its applications, vol. 17, Addison-Wesley, Reading (1983), 18–38.
- [15] M. Crochemore, “Tests sur les morphismes faiblement sans carré”, in: L. J. Cummings (ed.), *Combinatorics on Words*, Academic Press, Toronto (1983), 63–89.

- [16] R. O. Shelton, “On the Structure and Extendability of Square-Free Words”, in: L. J. Cummings (ed.), *Combinatorics on Words*, Academic Press, Toronto (1983), 101–118.
- [17] V. Elser, “Repeat-free sequences”, Lawrence Berkeley Laboratory preprint LBL-16632 (Sept. 1983).
- [18] F.-J. Brandenburg, “Uniformly growing k -th power-free homomorphisms”, *Theor. Comp. Sci.* **23** (1983) 69–82.
- [19] J. Brinkhuis, “Non-repetitive sequences on three symbols”, *Quart. J. Math. Oxford* **34** (1983) 145–149.
- [20] M. Leconte, “ k -th power-free codes”, in: M. Nivat and D. Perrin (eds.), *Automata on Infinite Words*, Lecture Notes in Computer Science 192, Springer, Berlin (1985), 172–187.
- [21] P. Séébold, “Overlap-free sequences”, in: M. Nivat and D. Perrin (eds.), *Automata on Infinite Words*, Lecture Notes in Computer Science 192, Springer, Berlin (1985), 207–215.
- [22] Y. Kobayashi, “Repetition-free words”, *Theor. Comp. Sci.* **44** (1986) 175–197.
- [23] P. A. B. Pleasants, private communication; and: “Entropies of sequences with the same language”, in preparation.